

A NEW SPECIES OF THE GENUS *TUAREGA* UVAROV FROM MOROCCO (ORTHOPTERA, PAMPHAGIDAE)

YIN Xiang-Chu^{1,2,3}, Martin Husemann⁴, LI Xin-Jiang^{1*}

1. College of Life Sciences, Hebei University, Baoding 071002, China; E-mail: yxch@sdu.edu.cn

2. Northwest Plateau Institute of Biology, Chinese Academy of Sciences, Xining 810008, China

3. College of Plant Protection, Shandong Agricultural University, Taian 271018, China

4. Biology Department, Baylor University, Waco, Texas 76706, USA; E-mail: Martin.Husemann@Baylor.edu

Abstract A new species, *Tuarega ouarzazatensis* sp. nov. (Orthoptera, Pamphagidae) is described from Morocco. The new species is similar to *Tuarega insignis* (Lucas, 1851), but differs from the latter in: the metazona of pronotum wider with widest part longer than length, median vein of tegmen not combined with cubital vein, the cubital vein of tegmen with 2 branches only. The type specimen is deposited in the Museum of Hebei University, China (MHU).

Key words Orthoptera, Pamphagidae, *Tuarega*, new species, Morocco.

The genus *Tuarega* was erected by Uvarov in 1943. The main characters of the genus are body large and robust; integument strongly rugose; fastigium of vertex short, wide, slightly sloping forwards and slightly concave, with fastigial furrow obliterated; antennae filiform, slightly shorter than head and pronotum together; pronotum in prozona almost cylindrical, metazona flatted, median carina weak, linear, crossed by three weak sulci, lateral carinae absent, metazona longer than prozona, its posterior margin elongated and angular, with obtuse apex; prosternum with very low collar; mesosternal interspace more than twice as wide as its length; tegmina and wings fully developed; tibia of second leg with a row of tubercles on upper side; hind femur moderately robust, with slightly expanded marginal areas; hind tibia with small apical spine on inner and outer side; arolium small; subtympal lobe large; Krauss' organ developed; first abdominal tergite above with large, ridge-like convexity. Male supra-anal plate elongate-angular; cerci narrow, incurved. Ovipositor moderately short, slender, with acute curved valves lower valve with large projection on the basal part. The type species is *Tuarega insignis* (Lucas, 1851) (= *Oedipoda insignis* Lucas, 1851). According to the Orthoptera species file online this is the only one species belong to this genus (Eades *et al.*, 2010). The type locality of the type species is Northern Africa, Algeria, Kefoum-Tebouc. *T. insignis* is distributed mainly in Northern Africa. Records from South Africa need to be verified (Lucas, 1851; Krauss, 1892; Kirby, 1910; Uvarov, 1943; Chopard, 1943, 1954;

Dekeyser and Villiers, 1956; Dirsh, 1956, 1965; Johnston, 1956, 1968; Korsakoff, 1958; Descamps, 1970; Yin, Shi and Yin, 1996; Massa and Rizzo, 1998; Usmani, 2007, 2008). In this paper a new species of the genus *Tuarega*, *Tuarega ouarzazatensis* sp. nov., is described from Ouarzazate of Morocco.

The type specimen is deposited in the Museum of Hebei University, China (MHU).

***Tuarega ouarzazatensis* sp. nov.** (Figs 1–6)


Holotype ♀, 10 km SE Ouarzazate, Morocco (30°51'N, 6°51'W; alt. 1 100 m), collected by MH and Dr. Jan Habel (Natural History Museum of Luxembourg) in 4 Mar. 2009.

Female. Body large and robust. Integument strongly rugose. Head subglobular, shorter than length of pronotum; fastigium of vertex short, wide, slightly sloping forwards and slightly concave, with fastigial furrow obliterated; frons vertical; frontal ridge low, above ocellus flat, at and below it concave, in basal part obliterated. Eyes oval, ratio of horizontal diameter: longitudinal diameter: subocular furrow as 3:4:5. Antennae filiform, 22 joints, slightly shorter than head and pronotum together. Pronotum in prozona almost cylindrical, metazona flattened; median carina weak, linear, crossed by three shallow indistinct sulci; lateral carinae absent; metazona longer than prozona; its posterior margin elongated and angular, with obtuse apex; width of metazona longer than its length. Prosternum with very low collar. Mesosternal interspace more than twice as wide as its length. Tegmina and wings fully developed, surpassing

This study was supported by the National Natural Sciences Foundation of China (30770263) and the Doctor Research Foundation of Hebei University (2009-167).

* Corresponding author, E-mail: hbulxj@yahoo.com.cn

Received 2 June 2010, accepted 8 June 2011.



Figs 1 – 6. *Tuarega ouarazatensis* sp. nov., female. 1. Lateral view. 2. Tibia of median leg. 3. Ovipositor valves. 4. Dorsal view. 5. Showing M and Cu veins. 6. Dorsal view of head and pronotum.

the end of abdomen distinctly. Cubital vein (Cu) of tegmina with 2 branches, distinct and not interconnected with median vein. Subtympanal lobe large. Krauss' organ developed. First abdominal tergite above with large, ridge-like convexity. Tibia of second leg with a row of tubercles on upper side. Hind femur moderately robust, with slightly expanded marginal areas. Hind tibia with small apical spine on inner and outer side, with 9 spines on upper margin of external and internal side (including apical spine). Arolium small. Cerci narrow, incurved, with obtuse

apex. Ovipositor moderately short, slender, with acute curved valves; lower valve with large projection on the basal part.

Coloration. Body brown. Tegmina yellowish-brown, with four pale dark bands on the V2 vein area and several dark brown spots. Hind wings yellow at base, with a dark strongly curved band reaching from the apical fourth to the posterior margin, but does not touch the outer margin, apical part with smoky spot, apical vein blackish. Hind femora brown. Hind tibiae (the specimen was deposited in the alcohol, so hind

tibiae changed into yellow on inner side, its original color was red) red on inner side. Ovipositor valves blackish.

Male. Unknown.

Measurements (in mm). Length of body 54.5; length of elytra 59.6; length of pronotum 16.3; length of hind femur 28.4.

Diagnosis. *Tuarega ouarazatensis* sp. nov. is similar to *Tuarega insignis* (Lucas, 1851), but differs

from the latter in three characters (Table 1): the metazona of pronotum wider, widest longer than length, median vein of tegmen not combined with cubital vein, the cubital vein of tegmen with 2 branches only.

Distribution. Currently only known from the type location.

Etymology. The specific epithet is named after the type locality, Ouarzazate, Morocco.

Table 1. Comparison of *Tuarega ouarazatensis* sp. nov. and *Tuarega insignis* (Lucas, 1851).

	<i>Tuarega ouarazatensis</i> sp. nov.	<i>Tuarega insignis</i> (Lucas, 1851)
Metazona of pronotum	Width longer than length	Width shorter than length
Tegmina	Median vein not combined with cubital vein	Median vein combined with cubital vein at posterior end
Cubital vein of tegmen	Having 2 branches	Having 3 branches

Acknowledgements We want to thank Dr. Jan Habel for organizing the collecting trips to Morocco and the Natural History Museum of Luxembourg for financing these field trips.

REFERENCES

Chopard, L. 1943. Orthoptéroïdes de L'Afrique du Nord. Faune de l'Empire Français, Lib. Larose, Paris. 1: 334
Chopard, L. 1954. Orthoptéroïdes de la région de Beni-Albes. *Bulletin de la Société Entomologique de France*, 59: 10–13.
Dirsh, V. M. 1956. The phallic complex in Acridoidea (Orthoptera) in relation to taxonomy. *Transactions of the Royal Entomological Society of London*, 108: 223–356, 66 pls.
Dirsh, V. M. 1965. The African genera of Acridoidea. Univ. Press, Cambridge. xiii + [i] + 578 + [2] pp., 452 figs.
Dekeyser, P. L. and Villiers, A. 1956. Contributions a letude du peuplement de la Mauretanie. Notations ecologiques et biogeographiques sur la faune de l'Adrar. *Mémoires de l'Institut Français d'Afrique Noire*, 44: 1–222, 25 pls.
Descamps, M. 1970. Contribution a la faune du Maroc, 3. Acridoidea du Maroc sararien et Dericorythinae (Orth.). *Bulletin de la Société Entomologique de France*, 75: 24–36, 6 figs.
Eades, D. C., Otte, D. and Naskrecki, P. 2010. Orthoptera Species File Online. Version 2.0/3.5. [16 Mar. 2010]. <http://osf2.orthoptera.org>.
Johnston, H. B. 1956. Annotated Catalogue of African Grasshoppers. The Cambridge University Press, Cambridge. 64.
Johnston, H. B. 1968. Annotated Catalogue of African Grasshoppers.

The Cambridge University Press, Cambridge. 62 (Suppl.): 448.
Kirby, W. F. 1910. A synonymic catalogue of Orthoptera (Orthoptera Saltatoria, Locustidae vel Acridiidae). *British Museum (Natural History)*, London, 3 (2): 290.
Korsakoff, M. N. 1958. Notes sur quelques insectes de Beni-Ounif. *Eos. Revista Española de Entomología (Eos)*, 34: 135–148, 27 figs.
Krauss, H. 1892. Dermapteren und Orthopteren aus Tunis. Gesammelt von Dr. A. König. *Wiener Entomologische Zeitung*, 11 (5): 149.
Lucas, H. 1851. Quelques remarques géographiques sur les Acridides qui habitent les possessions francaises due nord de l'Afrique. *Annales de la Société Entomologique de France*, 9 (2): 370.
Massa, B. and Rizzo, M. C. 1998. Orthoptera raccolti in Tunisia nel 19 secolo e conservati nel Museo Civico di Storia Naturale “G. Doria” di Genova [Orthoptera collected in Tunisia during the 19th century and preserved in the Museo Civico di Storia Naturale G. Doria, Genoa]. *Annali del Museo Civico di Storia Naturale ‘Giacomo Doria’*, Genova, 92: 280.
Usmani, 2007. Studies on some Libyan species of Pamphagidae (Orthoptera: Acridoidea). *Zootaxa*, 1625: 55.
Usmani, 2008. Studies on Acridoidea (Orthoptera) with some new records from Fezzan, Libya. *Zootaxa*, 1946: 13.
Uvarov, B. P. 1943. The tribe Thrinchini of the subfamily Pamphaginae, and the interrelations of the Acridid subfamilies Orthoptera). *Transactions of the Royal Entomological Society of London*, 93: 1–72, 73 figs.
Yin, X-C, Shi, J-P and Yin, Z 1996. A Synonymic Catalogue of Grasshoppers and Their Allies of the World (Orthoptera: Caelifera). China Forestry Publishing House, Beijing. 732.

摩洛哥土尔蝗属一新种（直翅目，癩蝗科）

印象初^{1,2,3} Martin Husemann⁴ 李新江^{1*}

1. 河北大学生命科学学院 保定 071002, E-mail: yxch@sda.u.edu.cn
2. 中国科学院西北高原生物研究所 西宁 810008
3. 山东农业大学植物保护学院 泰安 271018
4. Biology Department, Baylor University, Waco, Texas 76706, USA; E-mail: Martin_Husemann@Baylor.edu

摘 要 记述了采自摩洛哥土尔蝗属 *Tuarega* 1 新种, 瓦尔扎扎特土尔蝗 *Tuarega ouarazatensis* sp. nov., 模式标本保存于河北大学博物馆。

尔瓦扎扎特土尔蝗, 新种 *Tuarega ouarazatensis* sp. nov. (图 1~6)

新种与 *Tuarega insignis* (Lucas, 1851) 近似, 区别特征

* 通讯作者, E-mail: hbulxj@yahoo.com.cn

为：前胸背板沟后区宽大于长，前翅中脉与肘脉不闭合，前翅肘脉有 2 个分支。

正模 ♀，摩洛哥瓦尔扎扎特东南 10 km，海拔 1 100 m，2009-03-04，Martin Husemann 和 Jan Habel 博士（Natural

关键词 直翅目，癩蝗科，土尔蝗属，新种，摩洛哥.

中图分类号 Q969.26

History Museum of Luxembourg) 采。

词源：新种种名出自摩洛哥的瓦尔扎扎特，模式标本产地。